

Viewpoint

Improving Adherence to Behavioral Parent Training for ADHD Using Digital Health Tools

Linda J. Pfiffner^{1,*}, Melissa R. Dvorsky², Elizabeth J. Hawkey¹, Sara Chung¹,
Lauren M. Haack¹, Elizabeth B. Owens¹

¹ Department of Psychiatry and Behavioral Sciences, University of California San Francisco, San Francisco, CA 94143, USA

² Department of Pediatrics and Psychiatry and Behavioral Sciences, Children's National Hospital, The George Washington University School of Medicine and Health Sciences, Washington, DC 20010, USA

* Correspondence: Linda J. Pfiffner, Email: linda.pfiffner@ucsf.edu.

ABSTRACT

Behavioral Parent Training (BPT) is a well-established treatment for school-age children with ADHD but lack of parent adherence to prescribed parenting strategies limits treatment gains. Digital Health (dHealth) tools can be leveraged to target barriers to parent adherence but existing tools for parenting interventions are limited. New efforts to develop a dHealth tool to target adherence barriers including limited skill competence, EF processes, and low motivation/negative attitudes, are presented and recommendations for future technology-enhanced treatments are provided.

KEYWORDS: ADHD; behavioral parent training; digital health tools

IMPORTANCE OF PARENT ADHERENCE TO TREATMENT FOR ADHD

Parenting stress, parent-child conflict, and ineffective parenting are elevated in families of children with ADHD [1,2] and predict poor academic outcomes, interpersonal difficulties, and aggressive behavior [3,4]. BPT is a well-established ADHD treatment delivered in clinic settings [5] and more recently, school settings [6,7]. However, improvements in child behavior can be circumscribed and lack sustainability [8]. Poor parental adherence contributes to these limited effects [9,10], since BPT relies on parents (including primary caretakers) using recommended behavioral strategies with children on a regular basis in everyday contexts. Parents often have difficulty using prescribed strategies consistently, and further reduce their use once treatment ends [8]. Given estimates that 40%–60% of parents of children with ADHD have difficulty fully participating in treatment [8], and less than half of parents complete BPT homework assignments [11], suboptimal parent engagement and adherence are serious impediments to BPT effectiveness.

Open Access

Received: 14 July 2022

Accepted: 17 August 2022

Published: 22 August 2022

Copyright © 2022 by the author(s). Licensee Hapres, London, United Kingdom. This is an open access article distributed under the terms and conditions of [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

BARRIERS TO SKILL USE

Existing studies document barriers that can impede parent adherence including: (1) *limited skill competence* (e.g., inadequate understanding or execution of skills) [12,13], (2) *interfering executive functioning(EF) processes* (e.g., forgetting to use skills; inadequate prioritizing/planning; losing materials/resources) [1,13] and (3) *low motivation/negative attitudes* (e.g., low self-efficacy, pessimism, parenting stress) [1,12,14] as well as (4) *maintenance-specific barriers* when treatment ends (e.g., lack of success adapting learned skills to new challenges; decreased accountability). Notably, ADHD and EF-related problems are common among parents of youth with ADHD [15] and related adverse effects on parent adherence may be especially pronounced at follow-up [15,16].

Surprisingly little research has focused on reducing such barriers to BPT adherence, with most efforts focused on treatment access. Some behavioral treatments have begun to address skill implementation challenges secondary to parental ADHD symptoms by providing more flexible and individualized delivery [15]. However, these adaptations can be costly, unfeasible to deliver in limited-resourced school/community mental health settings and crucially, fail to address treatment sustainability [8,15]. Feasible and practical augmentations to BPTs that address adherence barriers and skill utilization are needed to improve sustained treatment outcomes.

ADDRESSING ADHERENCE BARRIERS WITH TECHNOLOGY

Use of technology to deliver or enhance mental health treatment has proliferated in recent years with smartphones providing on-demand and flexible access to treatment and thereby increasing treatment potency while minimizing costs [17–20]. Online portals and mobile applications that track adherence and provide feedback have demonstrated efficacy and feasibility for a variety of conditions, including for ADHD medication management [21,22].

Many digital tools specifically target parenting, but the vast majority are not empirically evaluated or based on evidence-based parenting practices [23]. Recently, however, several dHealth tools have been developed for early childhood BPTs and demonstrate promise either as augmentations [24] or alternatives to in-person treatment (“self-administered” programs) [25]. Nevertheless, existing BPT dHealth tools are limited. None have been designed for the specific needs related to parenting school-age youth with ADHD when BPT skills become more complex and children become more directly involved in treatment. Current solutions also do not specifically target adherence barriers faced by many parents of youth with ADHD who themselves are at increased risk for challenges with EF processes, low motivation, and parenting skill acquisition. Notably, use of existing technology is often less than optimal (e.g., 35%–50%+ fail to complete online modules [26,27]) and has not

incorporated features to support parenting in the period following active treatment when gains often diminish [28]. Digital tools are most likely to be used if they can be incorporated into daily routines and integrated into existing face-to-face treatments and may help to address contextual issues in the environment that affect adherence. For example, digital tools may be used to help parents track personalized daily goals for themselves and their children and can prompt parents when to use specific parenting skills in real-life contexts at prescribed times (e.g., morning routine, homework time). Such prompting could potentially include a location function that delivers contextually relevant guidance depending upon location (e.g., home, car, store, restaurant). Digital tools may also provide on-demand guidance for troubleshooting problematic situations via a digital wizard/coach through analysis of relevant antecedent and consequent events. However, tools have often been designed and developed with little input from the key stakeholders [29] (parents and children with ADHD, clinicians), resulting in products that may not be engaging or useful for families or clinicians and therefore less likely to facilitate adherence or sustainability. Conceptual frameworks for behavior change can guide optimization of the delivery interface in digital behavior change interventions, but these are often not applied during development. For example, psychological theories of persuasion and attitude change, motivation, and self-regulation can inform selection and design of features [30] which may better engage purported treatment mechanisms underlying behavior change and sustainability.

A NOVEL THEORY- AND STAKEHOLDER-INFORMED TOOL TO AUGMENT BPT

With support from the National Institute of Mental Health (R34 MH122222), our team is designing a dHealth tool to augment school-based BPT for ADHD that addresses the limitations above. Key stakeholders (parents, children, school clinicians) are informing each iteration of tool development in the context of a user-centered, mixed method design. As depicted in our conceptual model (Figure 1), the tool targets empirically-supported barriers to parent adherence [12,13]: skill competence, EF processes, and motivation/attitudes. We applied frameworks for digital behavior change interventions [30–32], to select the primary features addressing each barrier. For example, features addressing *Skill Competence* were informed by experiential learning theory [32] and include interactive practice activities via skill modules which provide opportunities for active learning and engage parents in setting their own personalized short-term goals and action planning [20,33]. Features for overcoming barriers due to *Executive Functioning Processes* are informed by self-regulation theories [30] and include automated reminders, streamlined content, and personalized goal-setting and action planning. Features to enhance *Motivation and Attitudinal Processes* for skill

utilization [34] are guided by social cognitive theory [30] and include personalized goal-setting, automated monitoring of parent/child progress, motivational prompts, and gamification (digital rewards for skill use). Linked parent/child views of child goals and reward plans integrated in daily routines address EF and motivational barriers since children who earn rewards when parents check their behavior goals are likely to remind and motivate parents to use the tool. To support sustained and independent parental skill use, *Maintenance Specific Features* include navigation for personalized problem-solving guides parents and troubleshooting new behavior problems by prompting learned skills. As depicted Figure 1, reductions in these targeted barriers are hypothesized to increase immediate and sustained parent skill utilization which in turn is expected to improve immediate and sustained child outcomes.




Figure 1. Theoretical Targets for Technology Solutions to Improve Parent Adherence and Sustained Outcomes in Behavior Parent Training for Youth with ADHD.

IMPLICATIONS FOR FUTURE RESEARCH

DHealth tools are promising, but have yet to be fully realized as effective methods for addressing adherence problems and sustaining treatment effects. Key elements likely to optimize effects of digital augmentations include: (1) developing tools specifically targeting empirically-supported barriers to adherence based on the intervention theory of change, (2) incorporating conceptually relevant and empirically-supported features, and (3) using an iterative user-centered design with families affected by the target condition and their clinicians to ensure usability and utility of the tool. The potential for broad uptake and scalability of digital interventions has never been greater. By including theory-informed features, targeting key treatment mechanisms, and involving stakeholders throughout the development process, current efforts leveraging technology will be better positioned to increase efficacy and reach of evidence-based practices.

CONFLICTS OF INTEREST

The authors declare that they have no conflicts of interest.

FUNDING

This research was funded by the National Institute of Mental Health, grant number R34 MH122222 and the University of California San Francisco Academic Senate (PI: Linda Pfiffner, PhD).

REFERENCES

1. Johnston C, Mash EJ. Families of children with attention-deficit/hyperactivity disorder: review and recommendations for future research. *Clin Child Fam Psychol Rev.* 2001;4(3):183-207.
2. Johnston C, Jassy JS. Attention-deficit/hyperactivity disorder and oppositional/conduct problems: Links to parent-child interactions. *J Can Acad Child Adolesc Psychiatry.* 2007;16(2):74.
3. Hinshaw SP, Owens EB, Wells KC, Kraemer HC, Abikoff HB, Arnold LE, et al. Family processes and treatment outcome in the MTA: Negative/ineffective parenting practices in relation to multimodal treatment. *J Abnorm Child Psychol.* 2000;28(6):555-68.
4. Molina BSG, Hinshaw SP, Swanson JM, Arnold LE, Vitiello B, Jensen PS, et al. The MTA at 8 years: prospective follow-up of children treated for combined-type ADHD in a multisite study. *J Am Acad Child Adolesc Psychiatry.* 2009;48(5):484-500.
5. Evans SW, Owens JS, Wymbs BT, Ray AR. Evidence-Based Psychosocial Treatments for Children and Adolescents With Attention Deficit/Hyperactivity Disorder. *J Clin Child Adolesc Psychol.* 2018;47(2):157-98. doi: 10.1080/15374416.2017.1390757
6. Fabiano GA, Schatz NK, Lupas K, Gordon C, Hayes T, Tower D, et al. A school-based parenting program for children with attention-deficit/hyperactivity disorder: Impact on paternal caregivers. *J Sch Psychol.* 2021;86(June):133-50. doi: 10.1016/j.jsp.2021.04.002
7. Pfiffner LJ, Rooney M, Haack L, Villodas M, Delucchi K, McBurnett K. A Randomized Controlled Trial of a School-Implemented School-Home Intervention for Attention-Deficit/Hyperactivity Disorder Symptoms and Impairment. *J Am Acad Child Adolesc Psychiatry.* 2016;55(9):762-70. doi: 10.1016/j.jaac.2016.05.023
8. Chacko A, Jensen SA, Lowry LS, Cornwell M, Chimklis A, Chan E, et al. Engagement in behavioral parent training: Review of the literature and implications for practice. *Clin Child Fam Psychol Rev.* 2016;19(3):204-15.
9. Clarke AT, Marshall SA, Mautone JA, Soffer SL, Jones HA, Costigan TE, et al. Parent Attendance and Homework Adherence Predict Response to a Family-School Intervention for Children With ADHD. *J Clin Child Adolesc Psychol.* 2015;44(1):58-67. doi: 10.1080/15374416.2013.794697
10. Dvorsky MR, Friedman LM, Spiess M, Pfiffner LJ. Patterns of Parental Adherence and the Association to Child and Parenting Outcomes Following a

- Multicomponent School-Home Intervention for Youth With ADHD. *Behav Ther.* 2021;52(3):745-60. doi: 10.1016/j.beth.2020.09.007
11. Haine-Schlagel R, Walsh NE. A Review of Parent Participation Engagement in Child and Family Mental Health Treatment. *Clin Child Fam Psychol Rev.* 2015;18(2):133-50. doi: 10.1007/s10567-015-0182-x
 12. Acri M, Chacko A, Gopalan G, McKay M. Engaging Families in Treatment for Child Behavior Disorders. *Wiley Handb Disruptive Impuls Disord.* 2017:391-409. doi: 10.1002/9781119092254.ch24
 13. Johnston C, Mah JWTT, Regambal M. Parenting Cognitions and Treatment Beliefs as Predictors of Experience Using Behavioral Parenting Strategies in Families of Children With Attention-Deficit/Hyperactivity Disorder. *Behav Ther.* 2010;41(4):491-504. doi: 10.1016/j.beth.2010.02.001
 14. Smith E, Koerting J, Latter S, Knowles MM, McCann DC, Thompson M, et al. Overcoming barriers to effective early parenting interventions for attention-deficit hyperactivity disorder (ADHD): Parent and practitioner views. *Child Care Health Dev.* 2015. doi: 10.1111/cch.12146
 15. Chronis-Tuscano A, Wang CH, Woods KE, Strickland J, Stein MA. Parent ADHD and evidence-based treatment for their children: review and directions for future research. *J Abnorm Child Psychol.* 2017;45(3):501-17.
 16. Dawson AE, Wymbs BT, Marshall SA, Mautone JA, Power TJ. The Role of Parental ADHD in Sustaining the Effects of a Family-School Intervention for ADHD. *J Clin Child Adolesc Psychol.* 2016;45(3):305-19. doi: 10.1080/15374416.2014.963858
 17. Jones DJ, Anton M, Gonzalez M, Honeycutt A, Khavjou O, Forehand R, et al. Incorporating mobile phone technologies to expand evidence-based care. *Cogn Behav Pract.* 2015;22(3):281-90.
 18. Jones DJ, Forehand R, Cuellar J, Parent J, Honeycutt A, Khavjou O, et al. Technology-Enhanced Program for Child Disruptive Behavior Disorders: Development and Pilot Randomized Control Trial. *J Clin Child Adolesc Psychol.* 2014;43(1):88-101. doi: 10.1080/15374416.2013.822308
 19. Breitenstein SM, Gross D, Christophersen R. Digital delivery methods of parenting training interventions: a systematic review. *Worldviews Evidence-Based Nurs.* 2014;11(3):168-76.
 20. Lindhiem O, Bennett CB, Orimoto TE, Kolko DJ. A meta-analysis of personalized treatment goals in psychotherapy: A preliminary report and call for more studies. *Clin Psychol Sci Pract.* 2016;23(2):165-76.
 21. Epstein JN, Kelleher KJ, Baum R, Brinkman WB, Peugh J, Gardner W, et al. Impact of a web-portal intervention on community ADHD care and outcomes. *Pediatrics.* 2016;138(2):e20154240.
 22. Wade SL, Walz NC, Carey J, Williams KM, Cass J, Herren L, et al. A randomized trial of teen online problem solving for improving executive function deficits following pediatric traumatic brain injury. *J Head Trauma Rehabil.* 2010;25(6):409-415.
 23. Baum RA, Epstein JN, Kelleher K. Healthcare reform, quality, and technology: ADHD as a case study. *Curr Psychiatry Rep.* 2013;15(7):369. doi: 10.1007/s11920-013-0369-0

24. Jones DJ, Forehand R, Cuellar J, Parent J, Honeycutt A, Khavjou O, et al. Technology-enhanced program for child disruptive behavior disorders: Development and pilot randomized control trial. *J Clin Child Adolesc Psychol*. 2014;43(1):88-101.
25. Baumel A, Pawar A, Mathur N, Kane JM, Correll CU. Technology-assisted parent training programs for children and adolescents with disruptive behaviors: A Systematic Review. *J Clin Psychiatry*. 2017;78(8):e957-69. doi: 10.4088/JCP.16r11063
26. Enebrink P, Högström J, Forster M, Ghaderi A. Internet-based parent management training: A randomized controlled study. *Behav Res Ther*. 2012 Apr;50(4):240-9. doi: 10.1016/j.brat.2012.01.006
27. Sanders MR, Baker S, Turner KMT. A randomized controlled trial evaluating the efficacy of Triple P Online with parents of children with early-onset conduct problems. *Behav Res Ther*. 2012 Nov;50(11):675-84. doi: 10.1016/j.brat.2012.07.004
28. Abikoff H, Gallagher R, Wells KC, Murray DW, Huang L, Lu F, et al. Remediating organizational functioning in children with ADHD: Immediate and long-term effects from a randomized controlled trial. *J Consult Clin Psychol*. 2013;81(1):113.
29. Mohr DC, Weingardt KR, Reddy M, Schueller SM. Three problems with current digital mental health research. and three things we can do about them. *Psychiatr Serv*. 2017;68(5):427-9. doi: 10.1176/appi.ps.201600541
30. Morrison LG. Theory-based strategies for enhancing the impact and usage of digital health behaviour change interventions: A review. *Digit Heal*. 2015;1:205520761559533. doi: 10.1177/2055207615595335
31. Hekler EB, Michie S, Pavel M, Rivera DE, Collins LM, Jimison HB, et al. Advancing Models and Theories for Digital Behavior Change Interventions. *Am J Prev Med*. 2016;51(5):825-32. doi: 10.1016/j.amepre.2016.06.013
32. Broussard JD, Teng EJ. Models for enhancing the development of experiential learning approaches within mobile health technologies. *Prof Psychol Res Pract*. 2019;50(3):195-203. doi: 10.1037/pro0000234
33. Lindhiem O, Kolko DJ, Cheng Y. Predicting psychotherapy benefit: A probabilistic and individualized approach. *Behav Ther*. 2012;43(2):381-92.
34. Cugelman B. Gamification: what it is and why it matters to digital health behavior change developers. *JMIR Serious Games*. 2013 Dec 12;1(1):e3.

How to cite this article:

Pfiffner LJ, Dvorsky MR, Hawkey EJ, Chung S, Haack LM, Owens EB Improving Adherence to Behavioral Parent Training for ADHD Using Digital Health Tools. *J Psychiatry Brain Sci*. 2022;7:e220005. <https://doi.org/10.20900/jpbs.20220005>