

*Mini Review*

## Astrocytes as Context for the Involvement of Myelin and Nodes of Ranvier in the Pathophysiology of Depression and Stress-Related Disorders

José Javier Miguel-Hidalgo

Department of Psychiatry and Human Behavior, University of Mississippi Medical Center, 2500 North State Street, Jackson, MS 39216, USA;  
Email: jmiguel-hidalgo@umc.edu.

---

### ABSTRACT

Astrocytes, despite some shared features as glial cells supporting neuronal function in gray and white matter, participate and adapt their morphology and neurochemistry in a plethora of distinct regulatory tasks in specific neural environments. In the white matter, a large proportion of the processes branching from the astrocytes' cell bodies establish contacts with oligodendrocytes and the myelin they form, while the tips of many astrocyte branches closely associate with nodes of Ranvier. Stability of myelin has been shown to greatly depend on astrocyte-to-oligodendrocyte communication, while the integrity of action potentials that regenerate at nodes of Ranvier has been shown to depend on extracellular matrix components heavily contributed by astrocytes. Several lines of evidence are starting to show that in human subjects with affective disorders and in animal models of chronic stress there are significant changes in myelin components, white matter astrocytes and nodes of Ranvier that have direct relevance to connectivity alterations in those disorders. Some of these changes involve the expression of connexins supporting astrocyte-to-oligodendrocyte gap junctions, extracellular matrix components produced by astrocytes around nodes of Ranvier, specific types of astrocyte glutamate transporters, and neurotrophic factors secreted by astrocytes that are involved in the development and plasticity of myelin. Future studies should further examine the mechanisms responsible for those changes in white matter astrocytes, their putative contribution to pathological connectivity in affective disorders, and the possibility of leveraging that knowledge to design new therapies for psychiatric disorders.

### Open Access

Received: 02 February 2023

Accepted: 21 February 2023

Published: 22 February 2023

Copyright © 2023 by the author(s). Licensee Hapres, London, United Kingdom. This is an open access article distributed under the terms and conditions of [Creative Commons Attribution 4.0 International License](#).

---

**KEYWORDS:** astrocytes; oligodendrocytes; depression; stress; myelin; connectivity; white matter

---

### INTRODUCTION

There is a solid body of experimental and human postmortem brain evidence demonstrating that astrocytes play fundamental roles in


supporting neuronal metabolism, survival and communication at synapses, and in the development and maintenance of the white matter. These essential roles and the ability of astrocytes to become reactive when confronted with injury or disease are reflected in their crucial involvement in the neuropathological mechanisms of most neurological and neurodegenerative disorders [1–3]. The recent decades have also witnessed increasing research attention towards the role of white matter pathology in brain connectivity changes associated with depression and other affective psychiatric disorders [4–7]. Connectivity involves the exchange of information between brain regions that is reflected in correlated functional changes [8,9] and also in structural alterations in the bundles of axons that actually propagate the signals to connect distant brain regions [10–12]. Part of the connectivity alterations in psychopathology can be accounted by local changes in synaptic transmission and cell metabolism in the gray matter (GM) [13–15]. These alterations will eventually result in undesired or maladaptive signal patterns that could simply be conveyed by axons in the WM to other brain destinations, so spreading dysfunction. In addition, regardless of neuronal, synaptic and glial [16] disturbances in the GM, the propagation of signals within the WM could be altered by pathology of the intrinsic WM components affecting the myelin insulation of axons or the structure and molecular composition of the axons themselves [17–20]. In fact, alterations intrinsic to the WM are now considered to be part of plastic processes occurring during learning and other normal non-pathological plastic processes in the brain [21]. Even in non-diseased brain structures, many studies have shown that the myelin that wraps many axons in the WM can undergo substantial morphological and neurochemical plastic changes that contribute to connectivity modifications leading to actual behavioral and cognitive adaptations, including significant changes in learning and memory [22,23]. Human postmortem studies in depression-diagnosed subjects and in animal models of potent risk factors for depression, such as stress, have also identified striking effects on the abundance and morphology of oligodendrocytes and myelin or their markers in some relevant brain regions [17,18,24–29]. Both in the gray and white matter several lines of research have shown that astrocytes appear to be seminal to basic cellular processes in neurons and myelin-forming oligodendrocytes that support connectivity [16,18,20,30,31]. In the following, there is a review of some major aspects of astrocyte neurochemistry and function that support those processes, with particular emphasis on astrocyte involvement in ensuring adequate conduction of nervous signals in the white matter.

### **GM ASTROCYTES INVOLVED IN PROCESSES AFFECTING CONNECTIVITY IN DEPRESSION AND STRESS**

Each astrocyte and its highly branched processes occupy a region of the gray matter that is adjacent to neighboring astrocytes territories with very

little overlap [32] (Figure 1), each astrocyte extending up to 200 µm in diameter in humans [33]. The linear reach of the processes is thus small relative to the reach of axons, although the influence of astrocyte cytoplasmic changes is somewhat extended by the gap junctions between astrocytes that permit cytoplasm to cytoplasm communications through shared transient intracellular calcium increases and the exchange of molecules no larger than 1.5 kdalton. Nevertheless, the reach and speed of these astrocyte-to-astrocyte communication channels are respectively limited and slow, so that long range connectivity is rather based in the long processes of neurons and their ability to speedily send action potentials that spearhead synaptic communication between widely separated brain regions. The contribution of astrocytes to long range communication does not depend thus on their individual reach within and among networks, but rather on the ability of astrocyte processes to regulate connectivity at the sites of neurotransmitter release, or locally support the metabolism of neurons through their ability to generate processes that branch out and produce leaflets closely apposed to other neural elements such as synapses or capillaries [34]. Even when glial cell communication may affect glial and neuronal activity in small local circuits, the main influence of astrocytes on longer range connections most likely depends on the function of astrocytes in reuptake of neurotransmitters glutamate and GABA, although other aspects of astrocyte function such as providing metabolic and neurotrophic substrates to neurons may also have an indirect permissive role for connectivity by sustaining ATP production and allowing dendritic and spine plasticity that are critical to maintenance and recovery of membrane potential and to synaptic release adaptations in neurons. The tips of the highly branched processes of astrocytes in the GM enwrap or are closely associated with presynaptic and their corresponding postsynaptic neuronal elements [34]. The pervasive presence of these trios of cellular elements (presynaptic-postsynaptic-astrocytic) in the gray matter, tightly undergirding synaptic communication, has given place to the concept of tripartite synapse. This arrangement of astrocyte processes allows singular astrocytes to affect as many as 100,000 synapses through the tips of their highly-branched processes [35,36]. In addition to influencing connectivity at the synaptic level, tips of astrocyte processes in the gray matter also closely appose nodes of Ranvier and the adjacent paranodal regions (Figure 1D) along myelinated portions of afferent and efferent axons that respectively come from or enter the white matter, so that part of the influence of gray matter astrocytes on connectivity may depend on their role at nodes of Ranvier in segments of axons residing in GM. However, the majority of astrocytes subserving NRs are expectedly located in the white matter [37], where they would exert critical permissive and regulatory roles for the propagation of action potentials [38], which are the signals carrying most of the information allowing for efficient mid- and long-range connectivity. Thus,

the involvement of WM astrocytes in processes influencing normal connectivity will be described and discussed in the next section.



**Figure 1.** Immunolabeled astrocytes in histological sections from the prefrontal cortical regions of rats (A, C, D) and from mixed-cell (astrocyte-neurons-oligodendrocytes) primary cultures from rat frontal cortex (B). Arrows in A, B and C point to the cell bodies of astrocytes. Red immunofluorescent label in A and B corresponds to astrocyte cytoskeletal marker GFAP. Cell nuclei in B are stained blue with nuclear marker DAPI. (C) GFAP labeling of astrocytes (blue) coexists with green label for glutamate transporter EAAT1 in cell bodies (arrow) and processes of astrocytes. D) White matter astrocytes display significant amounts of EAAT1, which was immunolabeled here to display blue fluorescence together with markers that identify nodes of Ranvier (beige arrows). Nodes of Ranvier are identified by using antibodies to neurofascin (red/pink) and to paranodal protein CASPR (green/yellow). Note that NRs are in close proximity to EAAT1-positive processes (blue immunofluorescence). “AS” denotes the nucleus of an EAAT1-positive astrocyte cell body in the white matter. Calibration bars at bottom left of each micrograph: (A) 50 μm, (B) 100 μm, (C) 15 μm, (D) 5 μm.

## WM ASTROCYTES INVOLVED IN PROCESSES AFFECTING CONNECTIVITY IN DEPRESSION AND STRESS

WM astrocytes are morphologically different from the large population of protoplasmic astrocytes in the GM, and are termed fibrous astrocytes [33,39]. Processes stemming from WM astrocytes are in overall thicker, fewer and less branched than those in the GM, although recent studies have shown and recognized the possibility of distinct morphological variation in WM astrocytes, particularly in response to WM injury [40]. In addition, unlike in GM, processes of WM astrocytes tend to intermingle in the WM without forming mutually exclusive territories [33]. Some of their branches abut the basal lamina around blood vessels to support the blood brain barrier and allow for regulated exchanges of water and metabolites with the blood circulation. In addition, many of the astrocytes' branches directly establish gap junctions with neighboring astrocytes and, importantly, with the cell membranes of oligodendrocytes. These gap junctions are formed by various types of connexins that are different in those between astrocytes as compared to those contacting astrocytes with oligodendrocytes. Various functions have been proposed for these cytoplasm-to-cytoplasm contacts, such as siphoning released extracellular potassium from axons to astrocytes, and eventually to the blood circulation, with the mediation of oligodendrocytes [41,42]. The critical role of gap junctions between astrocytes and oligodendrocytes to myelin maintenance is demonstrated in significant myelin disruptions observable in animal models deficient in one or more of the gap junction-forming connexins in astrocytes or oligodendrocytes [43–47]. Comparable alterations of astrocyte-oligodendrocyte gap junction-based communication may also contribute to the pathology of depression, since it has been found that gap junction coupling between astrocytes and oligodendrocytes may be reduced in the anterior cingulate cortex of subject with depression dying by suicide [48].

Altered structure and function of astrocytes and oligodendrocytes, and disturbed interactions between them have been proposed to play a major role in the pathophysiology of mood disorders, particularly MDD [49,50], while astrocyte atrophy is also being recently recognized to significantly occur during aging and neurodegeneration [51,52]. Low density of those cell types and reduced expression of astrocyte- and oligodendrocyte-specific gene transcripts were reported in postmortem human brains in MDD [17,19,20,53,54]. Similar features are observed in rodent models of depressive-like phenotypes caused by prolonged stress exposure such as chronic unpredictable stress [55–58]. However, most of these studies first focused on cortical gray matter and only later started to turn their attention to an involvement of white matter (WM) glial pathology as major contributor to functional and connectivity disturbances in MDD and models of stress-induced psychiatric disorders. Recently, prominent reductions in astrocyte density and astrocyte-specific protein and mRNA in ventral prefrontal WM in MDD [20,59] have been reported. In the same

brain region, oligodendrocyte size decreases and low myelin-specific gene expression was detected in WM of human subjects [17,19]. These findings align with reports of reduced fractional anisotropy (measure of axon bundles' integrity) in prefrontal WM [60] and corpus callosum [61] in MDD.

### ASTROCYTES AT NODES OF RANVIER

In addition to gap junction-mediated interactions with myelinated axons, the tips of a considerable proportion of WM astrocytic processes end in close apposition with nodes of Ranvier [38] (NRs), which are stretches of bare axon carrying ion channels for action potential regeneration and propagation. This closeness to NRs (Figure 1D) appears to be of great importance since more than 95% of NRs in WM are intimately associated with the ends of astrocytic processes (processes from another glial cell, NG2 cells, are also associated with NRs often in conjunction with astrocytes' processes) [37]. Astrocyte processes and partially unpacked portions of the oligodendrocyte-generated myelin coat around the axon (dubbed paranodes) are the main glial structures around NRs [37,62]. To stabilize the structure of NRs, and allow for focused aggregation of voltage gated at nodes of Ranvier, there are specialized cell-adhesion components of the oligodendrocyte membrane. These components face other adhesion components in the axonal membrane at each of the paranodes delimiting both ends of the NR [62,63]. Astrocytes have been proposed to regulate the speed of action potential propagation by influencing the separation between the paranodes surrounding the NR [38,63,64]. In addition, there is a distinct set of axonal membrane proteins anchored at the node itself, including neurofascin and contactin, that form a cell adhesion apparatus directly interacting with voltage-gated sodium channels [65,66]. At NRs, neurofascin 186, one of the cell adhesion proteins, in turn serves also as docking point for various extracellular matrix components such as proteoglycans and Tenascin-R, the precise functions of which are starting to be elucidated. Astrocytes provide a major contribution to these extracellular components, although oligodendrocytes, NG2 cells and the axons themselves contribute to the release of some of the particular proteoglycans or cell adhesion factors [67]. In fact, some of the extracellular proteoglycans such as proteoglycan BRAL-1 are heavily produced by astrocytes, and significant quantities of phosphacan, versican, brevican, neurocan and tenascin R are contributed by astrocytes as well [67–70].

Interaction between astrocyte processes and axons at NRs is mediated by those extracellular matrix (ECM) components, which include the proteoglycans brevican, phosphacan, versican-2, and BRAL-1, as well as the glycoprotein tenascin-R, and the axonal adhesion molecules neurofascin-186 (NF186) and contactin [62,71]. Despite detected alterations in astrocyte processes in GM and WM in depression much remains to be known about how the ubiquitous astrocyte processes at NRs

and the associated ECM molecules (ECMMs) and axon proteins are disturbed, and the role of those disturbances in connectivity changes. In a preliminary study in human PFC WM in MDD we found decreased levels of mRNAs for versican2, tenascin-R and NF186 and some of these alterations were still significant after removing outliers. Density of WM fibrous astrocytes has been found also decreased in CUS rats and in humans with MDD [20]. More recently we detected dramatic reductions in NR length and increases in the content of proteoglycan phosphacan associated with MDD and with chronic unpredictable stress in a rat model [72]. Thus, there appears to be significant pathology associated with WM astrocytes processes, astrocyte-linked ECMMs and axonal proteins at NRs. NR proteoglycans and tenascin-R bind to NR axon membrane proteins, forming a complex linking the axon cytoskeleton to the ECM. The link supports aggregation of axonal voltage-gated sodium channels ( $\text{Na}_v$ s) at NRs during NR formation and maintenance [73,74]. As stated above, astrocytes variably express all of these molecules [67]. Astrocyte processes matter to WM function because they are a major brevican, phosphacan and BRAL-1 source and these molecules allow for versican-2 and tenascin-R anchoring, which in turn bind to axonal adhesion proteins (NF186 and contactin) that regulate  $\text{Na}_v$ s' clustering [73]. Accordingly, proteoglycan-tenascin-R-NF186 complexes are thought crucial to support action potential regeneration at NRs for subsequent propagation. Thus, reduced or disrupted involvement of astrocytes processes in the generation of ECM components or in their display to support extracellular adhesion interactions may lead to alterations of signal propagation along myelinated axons and underlie a mechanism contributing to disturbed PFC connectivity in depression and other affective disorders.

## REGULATION OF GLUTAMATE ACTIONS IN THE WHITE MATTER

Although neuron-to-neuron synapses supporting neurotransmitter-based communication are mostly absent in the white matter, substantial glutamate release in the white matter has been detected [75]. This release is proposed to regulate the activity of glial cells, namely oligodendrocyte precursor cells (OPCs, also known as NG2 cells), since the remarkable discovery was made that synapses between presynaptic axons and postsynaptic OPC occur in the white matter [75], and that increased activity actually regulates the differentiations of OPC into oligodendrocytes to produce new myelin in a form of activity-dependent myelin plasticity [23]. WM astrocytes would play an important role in this type of plasticity because they express high levels of excitatory amino acid transporter 1 (EAAT1; in rodents the homologous protein is also called GLAST) glutamate transporters [37,76,77] (Figure 1C) that would facilitate reuptake of released glutamate and thus contribute to terminate the actions of WM extracellular glutamate, indirectly regulating the connectivity by controlling the activity of OPCs. Postmortem human research in the brains of MDD patients has detected a reduction on the

GFAP processes of astrocytes [20] and in size of CNPase-positive oligodendrocytes in the prefrontal white matter [78]. More recently low levels of GLAST proteins have been also measured in the ORB of human subjects with MDD [59] suggesting the possibility that a deficit of astrocyte processes or expression of EAAT1 may lead to a dysregulation of glutamate actions in the white matter.

Recent findings on the molecular composition of astrocyte processes suggest a role for ECMMs and EAAT1 transporters in regulating extracellular glutamate actions in the WM [71,79,80]. Both specific and cooperative actions of proteoglycans and EAAT1s of astrocytes at NRs contribute to extracellular glutamate regulation, NR maintenance, and stability of  $\text{Na}_v$  aggregation. These features are crucial for adequate propagation of action potentials. Thus, if glial processes, associated ECMMs and EAAT1s are altered in depression they could critically participate in pathophysiological mechanisms involving disturbances of connectivity among brain centers in MDD.

The processes of astrocytes associated with NRs, in conjunction with the myelin of oligodendrocytes, regulate the speed of saltatory conduction, thus allowing optimal communication between neurons [64,81]. Alterations of astrocyte processes abutting NRs and their vicinity could result in abnormal signal conduction and WM pathology as revealed by neuroimaging in depression [7,60,82–89]. Recently, abnormal structure and protein expression of axonal and oligodendrocyte proteins at NRs and adjacent axon segments were reported in corpus callosum and frontal cortex of chronically stressed rodents [61], suggesting that pathology at NRs may occur in MDD. So far it is unknown whether pathological alterations in astrocyte processes and NR-associated proteins in models of depression or in humans are direct participants in the etiology of depression. However, our previous research in astrocytes in MDD and our mRNA studies on WM would suggest that NR-related astrocytes changes contribute to depression-like dysfunction [17,72]. The possible involvement of astrocyte processes at NRs in WM functional and structural changes is consistent with reduced GFAP immunoreactive area fraction of processes and levels that we found in relatively young MDD subjects [53,90]. Since repeated stress is a risk factor for MDD [91] structural and molecular alterations associated with astrocyte processes at NRs may occur in prefrontal brain regions of MDD patients. Such pathology associated with astrocyte processes at NRs should be taken into account to explain the pathophysiology of depression or design innovative approaches for the treatment of affective and stress-related disorders.

## ASTROCYTES AND THE ROLE OF NEUROTROPHIC FACTORS

The importance of astrocytes to the maintenance of adequate signal conduction along axons also includes their role as providers of neurotrophic factors to facilitate oligodendrogenesis and myelination after damage to the white matter [92,93]. Other researchers have shown

that expression of neurotrophins by astrocytes, and their receptors in oligodendrocytes, is regulated by increased exercise [94], further suggesting that repair and plasticity of axons and their myelin is greatly influenced by neurotrophin release from astrocytes [93] even in non-pathological plastic processes. By contrast, in particular conditions involving inflammatory and toxic signals, and the mediation of neurotrophin receptors, dysfunctional astrocytes are potential enhancers of copper distribution and other deleterious compounds, and thus increase the possibility of myelin damage [93,95]. Other research has shown that myelination and the interaction between oligodendrocyte myelin membranes and axons at the paranodes lining NRs is dependent on astrocyte gene expression regulation by epigenetic factors such as Methyl-CpG-Binding Protein 2 (MeCP2), which in fact regulates the expression of growth factors such as BDNF and NGF by astrocytes [96]. The involvement of astrocyte-derived neurotrophic factors and their receptors in WM may be of great relevance to understand the contribution of neurotrophins to connectivity disturbances in depression since neurotrophic deficit or upregulation (depending on the brain area) have been found in the brains of affective disorder patients and in animal models of stress [97,98]. For example, significant correlation of neurotrophin DNA methylation and reduced WM integrity (assessed by neuroimaging methods) has been found in the WM of frontal brain regions in major depression [99]. Most importantly, the altered methylation patterns in depression subjects are strongly associated with astrocyte-related dysfunction [100]. Also in prefrontal brain regions, diffusion tensor magnetic resonance imaging (DTI) has shown increased fractional anisotropy in WM fiber bundles associated with particular genotypes of the neurotrophic tyrosine kinase receptor type 2 (*NTRK2*), but only in subjects diagnosed with depression [101], while specific polymorphisms of the BDNF gene are associated with detection of WM DTI disturbances in subjects with high depression severity [102,103].

## CONCLUDING REMARKS

Morphological and neurochemical differences of WM astrocytes as compared to most GM astrocytes, their ubiquitous involvement in myelin formation and maintenance, and their crucial association with nodes of Ranvier to support the regeneration of action potentials, suggest a critical contribution of those aspects of astrocyte biology to the integrity and fidelity of nervous signals carried along WM axons, and thus to normal brain connectivity. Accordingly, disturbances in brain connectivity that have been described in depression and other psychiatric disorders may be at least partially a consequence of disturbances in astrocyte interactions with oligodendrocyte membranes and nodes of Ranvier in the involved cortical brain regions. An argument could be made that targeting astrocyte-related pathology in the white matter may provide new or

complementary approaches to better therapies that improve connectivity in the brain of patients with affective and other psychiatric disorders.

## DATA AVAILABILITY

The dataset of the study is available from the authors upon reasonable request.

## CONFLICTS OF INTEREST

The author declares that he has no conflicts of interest.

## ACKNOWLEDGMENTS

Research reported in this article was supported by NIH grants R21MH118675 and R56MH113828.

## REFERENCES

1. Verkhratsky A, Zorec R, Parpura V. Stratification of astrocytes in healthy and diseased brain. *Brain Pathol.* 2017;27(5):629-44.
2. Escartin C, Galea E, Lakatos A, O'Callaghan JP, Petzold GC, Serrano-Pozo A, et al. Reactive astrocyte nomenclature, definitions, and future directions. *Nat Neurosci.* 2021;24(3):312-25.
3. Pekny M, Pekna M, Messing A, Steinhauser C, Lee JM, Parpura V, et al. Astrocytes: a central element in neurological diseases. *Acta Neuropathol.* 2016;131(3):323-45.
4. Kumar A, Cook IA. White matter injury, neural connectivity and the pathophysiology of psychiatric disorders. *Dev Neurosci.* 2002;24(4):255-61.
5. Zheng KZ, Wang HN, Liu J, Xi YB, Li L, Zhang X, et al. Incapacity to control emotion in major depression may arise from disrupted white matter integrity and OFC-amygdala inhibition. *CNS Neurosci Ther.* 2018;24(11):1053-62.
6. De Witte NAJ, Mueller SC. White matter integrity in brain networks relevant to anxiety and depression: evidence from the human connectome project dataset. *Brain Imaging Behav.* 2017;11(6):1604-15.
7. Jalbrzikowski M, Larsen B, Hallquist MN, Foran W, Calabro F, Luna B. Development of White Matter Microstructure and Intrinsic Functional Connectivity Between the Amygdala and Ventromedial Prefrontal Cortex: Associations With Anxiety and Depression. *Biol Psychiatry.* 2017;82(7):511-21.
8. Kaiser RH, Andrews-Hanna JR, Wager TD, Pizzagalli DA. Large-Scale Network Dysfunction in Major Depressive Disorder: A Meta-analysis of Resting-State Functional Connectivity. *JAMA Psychiatry.* 2015;72(6):603-11.
9. Rolls ET, Cheng W, Gilson M, Qiu J, Hu Z, Ruan H, et al. Effective Connectivity in Depression. *Biol Psychiatry Cogn Neurosci Neuroimaging.* 2018;3(2):187-97.
10. Drevets WC, Price JL, Furey ML. Brain structural and functional abnormalities in mood disorders: implications for neurocircuitry models of depression. *Brain Struct Funct.* 2008;213(1-2):93-118.

11. Liu J, Xu X, Zhu C, Luo L, Wang Q, Xiao B, et al. Disrupted Structural Brain Network Organization Behind Depressive Symptoms in Major Depressive Disorder. *Front Psychiatry*. 2020;11:565890.
12. de Kwaasteniet B, Ruhe E, Caan M, Rive M, Olabarriaga S, Groefsema M, et al. Relation between structural and functional connectivity in major depressive disorder. *Biol Psychiatry*. 2013;74(1):40-7.
13. Duman RS, Aghajanian GK, Sanacora G, Krystal JH. Synaptic plasticity and depression: new insights from stress and rapid-acting antidepressants. *Nat Med*. 2016;22(3):238-49.
14. Christoffel DJ, Golden SA, Russo SJ. Structural and synaptic plasticity in stress-related disorders. *Rev Neurosci*. 2011;22(5):535-49.
15. Vose LR, Stanton PK. Synaptic Plasticity, Metaplasticity and Depression. *Curr Neuropharmacol*. 2017;15(1):71-86.
16. Miguel-Hidalgo JJ. Astroglia in the Vulnerability to and Maintenance of Stress-Mediated Neuropathology and Depression. *Front Cell Neurosci*. 2022;16:869779.
17. Miguel-Hidalgo JJ, Hall KO, Bonner H, Roller AM, Syed M, Park CJ, et al. MicroRNA-21: Expression in oligodendrocytes and correlation with low myelin mRNAs in depression and alcoholism. *Prog Neuropsychopharmacol Biol Psychiatry*. 2017;79(Pt B):503-14.
18. Miguel-Hidalgo JJ, Moulana M, Deloach PH, Rajkowska G. Chronic Unpredictable Stress Reduces Immunostaining for Connexins 43 and 30 and Myelin Basic Protein in the Rat Prelimbic and Orbitofrontal Cortices. *Chronic Stress*. 2018;2:1-12.
19. Rajkowska G, Mahajan G, Maciag D, Sathyanesan M, Iyo AH, Moulana M, et al. Oligodendrocyte morphometry and expression of myelin-related mRNA in ventral prefrontal white matter in major depressive disorder. *J Psychiatr Res*. 2015;65:53-62.
20. Rajkowska G, Legutko B, Moulana M, Syed M, Romero DG, Stockmeier CA, et al. Astrocyte pathology in the ventral prefrontal white matter in depression. *J Psychiatr Res*. 2018;102:150-8.
21. Fields RD. White matter in learning, cognition and psychiatric disorders. *Trends Neurosci*. 2008;31(7):361-70.
22. Fields RD, Woo Dong H, Bassar Peter J. Glial Regulation of the Neuronal Connectome through Local and Long-Distant Communication. *Neuron*. 2015;86(2):374-86.
23. Fields RD. A new mechanism of nervous system plasticity: activity-dependent myelination. *Nat Rev Neurosci*. 2015;16(12):756-67.
24. Gao Y, Ma J, Tang J, Liang X, Huang CX, Wang SR, et al. White matter atrophy and myelinated fiber disruption in a rat model of depression. *J Comp Neurol*. 2017;525(8):1922-33.
25. Lehmann ML, Weigel TK, Elkahloun AG, Herkenham M. Chronic social defeat reduces myelination in the mouse medial prefrontal cortex. *Sci Rep*. 2017;7:46548.

26. Liu J, Dietz K, Hodes GE, Russo SJ, Casaccia P. Widespread transcriptional alternations in oligodendrocytes in the adult mouse brain following chronic stress. *Dev Neurobiol.* 2018;78(2):152-62.
27. Liu J, Dietz K, Deloyht JM, Pedre X, Kelkar D, Kaur J, et al. Impaired adult myelination in the prefrontal cortex of socially isolated mice. *Nat Neurosci.* 2012;15:1621-3.
28. Antontseva E, Bondar N, Reshetnikov V, Merkulova T. The Effects of Chronic Stress on Brain Myelination in Humans and in Various Rodent Models. *Neuroscience.* 2020;441:226-38.
29. Bonnefil V, Dietz K, Amatruda M, Wentling M, Aubry AV, Dupree JL, et al. Region-specific myelin differences define behavioral consequences of chronic social defeat stress in mice. *Elife.* 2019;8:e40855.
30. Lou YX, Li J, Wang ZZ, Xia CY, Chen NH. Glucocorticoid receptor activation induces decrease of hippocampal astrocyte number in rats. *Psychopharmacology (Berl).* 2018;235(9):2529-40.
31. Murphy-Royal C, Gordon GR, Bains JS. Stress-induced structural and functional modifications of astrocytes-Further implicating glia in the central response to stress. *Glia.* 2019;67(10):1806-20.
32. Sofroniew MV, Vinters HV. Astrocytes: biology and pathology. *Acta Neuropathol.* 2010;119(1):7-35.
33. Oberheim NA, Takano T, Han X, He W, Lin JH, Wang F, et al. Uniquely hominid features of adult human astrocytes. *J Neurosci.* 2009;29(10):3276-87.
34. Semyanov A, Verkhratsky A. Astrocytic processes: from tripartite synapses to the active milieu. *Trends Neurosci.* 2021;44(10):781-92.
35. Halassa MM, Fellin T, Takano H, Dong JH, Haydon PG. Synaptic islands defined by the territory of a single astrocyte. *J Neurosci.* 2007;27(24):6473-7.
36. Ogata K, Kosaka T. Structural and quantitative analysis of astrocytes in the mouse hippocampus. *Neuroscience.* 2002;113(1):221-33.
37. Serwanski DR, Jukkola P, Nishiyama A. Heterogeneity of astrocyte and NG2 cell insertion at the node of ranvier. *J Comp Neurol.* 2017;525(3):535-52.
38. Butt AM, Verkhratsky A. Astrocytes regulate action potential propagation in myelinated axons: It is very crowded at the node of Ranvier. *Cell Calcium.* 2022;101:102518.
39. Miller RH, Raff MC. Fibrous and protoplasmic astrocytes are biochemically and developmentally distinct. *J Neurosci.* 1984;4(2):585-92.
40. Bugiani M, Plug BC, Man JHK, Breur M, van der Knaap MS. Heterogeneity of white matter astrocytes in the human brain. *Acta Neuropathol.* 2022;143(2):159-77.
41. Kamasawa N, Sik A, Morita M, Yasumura T, Davidson KG, Nagy JI, et al. Connexin-47 and connexin-32 in gap junctions of oligodendrocyte somata, myelin sheaths, paranodal loops and Schmidt-Lanterman incisures: implications for ionic homeostasis and potassium siphoning. *Neuroscience.* 2005;136(1):65-86.
42. Rash JE. Molecular disruptions of the pangial syncytium block potassium siphoning and axonal saltatory conduction: pertinence to neuromyelitis

- optica and other demyelinating diseases of the central nervous system. *Neuroscience*. 2010;168(4):982-1008.
43. Lutz SE, Zhao Y, Gulinello M, Lee SC, Raine CS, Brosnan CF. Deletion of astrocyte connexins 43 and 30 leads to a dysmyelinating phenotype and hippocampal CA1 vacuolation. *J Neurosci*. 2009;29(24):7743-52.
  44. Sargiannidou I, Markoullis K, Kleopa KA. Molecular mechanisms of gap junction mutations in myelinating cells. *Histol Histopathol*. 2010;25(9):1191-206.
  45. Gankam Kengne F, Nicaise C, Soupart A, Boom A, Schiettecatte J, Pochet R, et al. Astrocytes are an early target in osmotic demyelination syndrome. *J Am Soc Nephrol*. 2011;22(10):1834-45.
  46. Cotrina ML, Nedergaard M. Brain connexins in demyelinating diseases: Therapeutic potential of glial targets. *Brain Res*. 2012;1487:61-8.
  47. Li T, Giaume C, Xiao L. Connexins-Mediated Glia Networking Impacts Myelination and Remyelination in the Central Nervous System. *Molecular Neurobiology*. 2014;49(3):1460-71.
  48. Tanti A, Lutz PE, Kim J, O'Leary L, Theroux JF, Turecki G, et al. Evidence of decreased gap junction coupling between astrocytes and oligodendrocytes in the anterior cingulate cortex of depressed suicides. *Neuropsychopharmacology*. 2019 Nov;44(12):2099-2111.
  49. Rajkowska G, Miguel-Hidalgo JJ. Gliogenesis and glial pathology in depression. *CNS Neurol Disord Drug Targets*. 2007;6(3):219-33.
  50. Wang Q, Timberlake II MA, Prall K, Dwivedi Y. The recent progress in animal models of depression. *Prog Neuropsychopharmacol Biol Psychiatry*. 2017 Jul 3;77:99-109.
  51. Verkhratsky A, Rodrigues JJ, Pivoriunas A, Zorec R, Semenyanov A. Astroglial atrophy in Alzheimer's disease. *Pflugers Arch*. 2019;471(10):1247-61.
  52. Verkhratsky A, Augusto-Oliveira M, Pivoriunas A, Popov A, Brazhe A, Semenyanov A. Astroglial asthenia and loss of function, rather than reactivity, contribute to the ageing of the brain. *Pflugers Arch*. 2021;473(5):753-74.
  53. Miguel-Hidalgo JJ, Baucom C, Dilley G, Overholser JC, Meltzer HY, Stockmeier CA, et al. Glial fibrillary acidic protein immunoreactivity in the prefrontal cortex distinguishes younger from older adults in major depressive disorder. *Biol Psychiatry*. 2000;48(8):861-73.
  54. Chandley MJ, Szebeni K, Szebeni A, Crawford J, Stockmeier CA, Turecki G, et al. Gene expression deficits in pontine locus coeruleus astrocytes in men with major depressive disorder. *J Psychiatry Neurosci*. 2013;38(4):276-84.
  55. Banasr M, Chowdhury GM, Terwilliger R, Newton SS, Duman RS, Behar KL, et al. Glial pathology in an animal model of depression: reversal of stress-induced cellular, metabolic and behavioral deficits by the glutamate-modulating drug riluzole. *Mol Psychiatry*. 2010;15(5):501-11.
  56. Heine VM, Maslam S, Zareno J, Joels M, Lucassen PJ. Suppressed proliferation and apoptotic changes in the rat dentate gyrus after acute and chronic stress are reversible. *Eur J Neurosci*. 2004;19(1):131-44.

57. Nollet M, Le Guisquet AM, Belzung C. Models of depression: unpredictable chronic mild stress in mice. *Curr Protoc Pharmacol.* 2021 Aug;1(8):e208. doi: 10.1002/cpz1.208.
58. Sun JD, Liu Y, Yuan YH, Li J, Chen NH. Gap junction dysfunction in the prefrontal cortex induces depressive-like behaviors in rats. *Neuropsychopharmacology.* 2012;37(5):1305-20.
59. Miguel-Hidalgo JJ, Waltzer R, Whittom AA, Austin MC, Rajkowska G, Stockmeier CA. Glial and glutamatergic markers in depression, alcoholism, and their comorbidity. *J Affect Disord.* 2010;127(1-3):230-40.
60. Tham MW, Woon PS, Sum MY, Lee TS, Sim K. White matter abnormalities in major depression: evidence from post-mortem, neuroimaging and genetic studies. *J Affect Disord.* 2011;132(1-2):26-36.
61. Miyata S, Taniguchi M, Koyama Y, Shimizu S, Tanaka T, Yasuno F, et al. Association between chronic stress-induced structural abnormalities in Ranvier nodes and reduced oligodendrocyte activity in major depression. *Sci Rep.* 2016;6:23084.
62. Rasband MN, Peles E. The Nodes of Ranvier: Molecular Assembly and Maintenance. *Cold Spring Harb Perspect Biol.* 2015;8(3):a020495.
63. Rasband MN, Peles E. Mechanisms of node of Ranvier assembly. *Nat Rev Neurosci.* 2021;22(1):7-20.
64. Dutta DJ, Woo DH, Lee PR, Pajevic S, Bukalo O, Huffman WC, et al. Regulation of myelin structure and conduction velocity by perinodal astrocytes. *Proc Natl Acad Sci U S A.* 2018;115(46):11832-7.
65. Lustig M, Zanazzi G, Sakurai T, Blanco C, Levinson SR, Lambert S, et al. Nr-CAM and neurofascin interactions regulate ankyrin G and sodium channel clustering at the node of Ranvier. *Current Biol.* 2001;11(23):1864-9.
66. Falk J, Bonnon C, Girault J-A, Faivre-Sarrailh C. F3/contactin, a neuronal cell adhesion molecule implicated in axogenesis and myelination. *Biol Cell.* 2002;94(6):327-34.
67. Song I, Dityatev A. Crosstalk between glia, extracellular matrix and neurons. *Brain Res Bull.* 2018;136:101-8.
68. Bekku Y, Vargova L, Goto Y, Vorisek I, Dmytrenko L, Narasaki M, et al. Bral1: its role in diffusion barrier formation and conduction velocity in the CNS. *J Neurosci.* 2010;30(8):3113-23.
69. Cicanic M, Sykova E, Vargova L. Bral1: “Superglue” for the extracellular matrix in the brain white matter. *Int J Biochem Cell Biol.* 2012;44(4):596-9.
70. Tanti A, Belliveau C, Nagy C, Maitra M, Denux F, Perlman K, et al. Child abuse associates with increased recruitment of perineuronal nets in the ventromedial prefrontal cortex: a possible implication of oligodendrocyte progenitor cells. *Mol Psychiatry.* 2022;27(3):1552-61.
71. Faivre-Sarrailh C, Devaux JJ. Neuro-glial interactions at the nodes of Ranvier: implication in health and diseases. *Front Cell Neurosci.* 2013;7:196.
72. Miguel-Hidalgo JJ, Hearn E, Holmes M, Clark A, Saleem K, Rajkowska G. Pathology of Nodes of Ranvier in the Prefrontal Cortex is Comparable in Depressed Subjects and Rats Under Chronic Stress. *Biol Psychiatry.* 2021;89(9):S204.

73. Nelson AD, Jenkins PM. Axonal Membranes and Their Domains: Assembly and Function of the Axon Initial Segment and Node of Ranvier. *Front Cell Neurosci.* 2017;11:136.
74. Susuki K, Chang KJ, Zollinger DR, Liu Y, Ogawa Y, Eshed-Eisenbach Y, et al. Three mechanisms assemble central nervous system nodes of Ranvier. *Neuron.* 2013;78(3):469-82.
75. Kukley M, Capetillo-Zarate E, Dietrich D. Vesicular glutamate release from axons in white matter. *Nat Neurosci.* 2007;10(3):311-20.
76. Douyard J, Shen L, Huganir RL, Rubio ME. Differential neuronal and glial expression of GluR1 AMPA receptor subunit and the scaffolding proteins SAP97 and 4.1N during rat cerebellar development. *J Comp Neurol.* 2007;502(1):141-56.
77. Chung EK, Chen LW, Chan YS, Yung KK. Downregulation of glial glutamate transporters after dopamine denervation in the striatum of 6-hydroxydopamine-lesioned rats. *J Comp Neurol.* 2008;511(4):421-37.
78. Miguel-Hidalgo JJ, Waltzer R, Lo V, Stockmeier C. Density of Cnpase-Positive Cells and Cnpase Levels in the Orbitofrontal Cortex in Alcoholism and Depression. *Alcohol Clin Exp Res.* 2009;33(6):26A.
79. Okuda H, Tatsumi K, Morita S, Shibukawa Y, Korekane H, Horii-Hayashi N, et al. Chondroitin sulfate proteoglycan tenascin-R regulates glutamate uptake by adult brain astrocytes. *J Biol Chem.* 2014;289(5):2620-31.
80. Sirko S, von Holst A, Weber A, Wizenmann A, Theocharidis U, Gotz M, et al. Chondroitin sulfates are required for fibroblast growth factor-2-dependent proliferation and maintenance in neural stem cells and for epidermal growth factor-dependent migration of their progeny. *Stem Cells.* 2010;28(4):775-87.
81. Stadelmann C, Timmler S, Barrantes-Freer A, Simons M. Myelin in the Central Nervous System: Structure, Function, and Pathology. *Physiol Rev.* 2019;99(3):1381-431.
82. Murphy ML, Frodl T. Meta-analysis of diffusion tensor imaging studies shows altered fractional anisotropy occurring in distinct brain areas in association with depression. *Biol Mood Anxiety Disord.* 2011;1(1):3.
83. Nobuhara K, Okugawa G, Sugimoto T, Minami T, Tamagaki C, Takase K, et al. Frontal white matter anisotropy and symptom severity of late-life depression: a magnetic resonance diffusion tensor imaging study. *J Neurol Neurosurg Psychiatry.* 2006;77(1):120-2.
84. Sexton CE, Mackay CE, Ebmeier KP. A systematic review of diffusion tensor imaging studies in affective disorders. *Biol Psychiatry.* 2009;66(9):814-23.
85. Shimony JS, Sheline YI, D'Angelo G, Epstein AA, Benzingher TL, Mintun MA, et al. Diffuse microstructural abnormalities of normal-appearing white matter in late life depression: a diffusion tensor imaging study. *Biol Psychiatry.* 2009;66(3):245-52.
86. Zheng C, Quan M, Zhang T. Decreased thalamo-cortical connectivity by alteration of neural information flow in theta oscillation in depression-model rats. *J Comput Neurosci.* 2012;33(3):547-58.

87. Zhu C, Ou L, Geng Q, Zhang M, Ye R, Chen J, et al. Association of somatic symptoms with depression and anxiety in clinical patients of general hospitals in Guangzhou, China. *Gen Hosp Psychiatry.* 2012;34(2):113-20.
88. Korgaonkar MS, Cooper NJ, Williams LM, Grieve SM. Mapping inter-regional connectivity of the entire cortex to characterize major depressive disorder: a whole-brain diffusion tensor imaging tractography study. *Neuroreport.* 2012;23(9):566-71.
89. Scheinost D, Holmes SE, DellaGioia N, Schleifer C, Matuskey D, Abdallah CG, et al. Multimodal Investigation of Network Level Effects Using Intrinsic Functional Connectivity, Anatomical Covariance, and Structure-to-Function Correlations in Unmedicated Major Depressive Disorder. *Neuropsychopharmacology.* 2018 Apr;43(5):1119-27.
90. Si XH, Miguel-Hidalgo JJ, O'Dwyer G, Stockmeier CA, Rajkowska G. Age-dependent reductions in the level of glial fibrillary acidic protein in the prefrontal cortex in major depression. *Neuropsychopharmacology.* 2004;29(11):2088-96.
91. Lupien SJ, McEwen BS, Gunnar MR, Heim C. Effects of stress throughout the lifespan on the brain, behaviour and cognition. *Nat Rev Neurosci.* 2009;10(6):434-45.
92. Miyamoto N, Maki T, Shindo A, Liang AC, Maeda M, Egawa N, et al. Astrocytes Promote Oligodendrogenesis after White Matter Damage via Brain-Derived Neurotrophic Factor. *J Neurosci.* 2015;35(41):14002-8.
93. Lundgaard I, Osorio MJ, Kress BT, Sanggaard S, Nedergaard M. White matter astrocytes in health and disease. *Neuroscience.* 2014;276:161-73.
94. Skup M, Dwornik A, Macias M, Sulejczak D, Wiater M, Czarkowska-Bauch J. Long-term locomotor training up-regulates TrkB(FL) receptor-like proteins, brain-derived neurotrophic factor, and neurotrophin 4 with different topographies of expression in oligodendroglia and neurons in the spinal cord. *Exp Neurol.* 2002;176(2):289-307.
95. Colombo E, Triolo D, Bassani C, Bedogni F, Di Dario M, Dina G, et al. Dysregulated copper transport in multiple sclerosis may cause demyelination via astrocytes. *Proc Natl Acad Sci U S A.* 2021;118(27):e2025804118.
96. Buch L, Lipi B, Langhnoja J, Jaldeep L, Pillai PP, Prakash P. Role of astrocytic MeCP2 in regulation of CNS myelination by affecting oligodendrocyte and neuronal physiology and axo-glial interactions. *Exp Brain Res.* 2018;236(11):3015-27.
97. Duman RS. Neurotrophic Factors in Etiology and Treatment of Mood Disorders. *Handbook of Contemporary Neuropharmacology.* New York (US): John Wiley & Sons, Inc.; 2007.
98. Castrén E, Vöikar V, Rantamäki T. Role of neurotrophic factors in depression. *Curr Opin Pharmacol.* 2007;7(1):18-21.
99. Choi S, Han KM, Won E, Yoon BJ, Lee MS, Ham BJ. Association of brain-derived neurotrophic factor DNA methylation and reduced white matter integrity in the anterior corona radiata in major depression. *J Affect Disord.* 2015;172:74-80.

100. Nagy C, Suderman M, Yang J, Szyf M, Mechawar N, Ernst C, et al. Astrocytic abnormalities and global DNA methylation patterns in depression and suicide. *Mol Psychiatry*. 2015;20(3):320-8.
101. Murphy ML, Carballido A, Fagan AJ, Morris D, Fahey C, Meaney J, et al. Neurotrophic tyrosine kinase polymorphism impacts white matter connections in patients with major depressive disorder. *Biol Psychiatry*. 2012;72(8):663-70.
102. Tatham EL, Ramasubbu R, Gaxiola-Valdez I, Cortese F, Clark D, Goodyear B, et al. White matter integrity in major depressive disorder: Implications of childhood trauma, 5-HTTLPR and BDNF polymorphisms. *Psychiatry Res Neuroimaging*. 2016;253:15-25.
103. Carballido A, Amico F, Ugwu I, Fagan AJ, Fahey C, Morris D, et al. Reduced fractional anisotropy in the uncinate fasciculus in patients with major depression carrying the met-allele of the Val66Met brain-derived neurotrophic factor genotype. *Am J Med Genet B*. 2012;159B(5):537-48.

How to cite this article:

Miguel-Hidalgo JJ. Astrocytes as Context for the Involvement of Myelin and Nodes of Ranvier in the Pathophysiology of Depression and Stress-Related Disorders. *J Psychiatry Brain Sci.* 2023;8:e230001.  
<https://doi.org/10.20900/jpbs.20230001>